Homogeneous nucleation and its role in CRYSTAL-FACE anvil cirrus— Part II: Model results

Ann Fridlind, Andy Ackerman, and Eric Jensen

NASA Ames Research Center

Dave Stevens

Lawrence Livermore National Laboratory

Donghai Wang

NASA Langley Research Center

Andy Heymsfield and Larry Miloshevich

National Center for Atmospheric Research

Cindy Twohy

Oregon State University

Mike Poellet


University of North Dakota


Tim VanReken, Tracey Rissman, Varuntida Varutbangkul, Rick Flagan, and John Seinfeld
California Institute of Technology


Haf JonssonCIRPAS


Motivation


- Anvil crystals are formed in strong updraft cores
 - few observations
 - high experimental uncertainty
 - complex microphysical environment
- Can homogeneous nucleation produce most ice crystals?
- Are boundary layer or free tropospheric aerosols more important?
- Is Hallett-Mossop ice multiplication important?
- Is heterogeneous ice nucleation important?



















DHARMA model description


- Domain resolution
 - □ 48 x 48 x 24 km
 - \Box dx = dy = 500 m, dz = 375 m
- Large-eddy simulation dynamics, dt = 5 s
 - open boundary conditions
 - assimilated meteorology (Miami sounding)
 - assimilated surface fluxes (ARPS model)
- Explicit microphysics, minimum dt = 0.2 s
 - □ 16 bins: aerosols (20 nm -1 um), liquid and ice (2 um ->1cm)
 - aerosol activation/freezing, condensation/evaporation, coagulation, breakup, homogeneous/heterogeneous ice nucleation, Hallett-Mossop ice multiplication


Assimilated surface fluxes: ARPS model





Assimilated aerosol profiles: Twin Otter and Citation


























Citation flight data






Can homogeneous nucleation produce most ice crystals?


Observations versus model layer at T = -36C





Observations versus model layer at T = -36C





Observations versus model layer at T = -36C















Boundary layer aerosol increased to 6500/mL




3.0


Boundary layer aerosol reduced to 400/mL




FT aerosol reduced to 10/mL above 2 km




FT aerosol reduced to 10/mL above 4 km




FT aerosol reduced to 10/mL above 6 km





FT aerosol reduced to 10/mL above 8 km







Hallett-Mossop ice crystal multiplication










30/L ice nuclei active at -25C





3000/L ice nuclei active at -25C





Preliminary conclusions for July 18

- can homogeneous nucleation produce most ice crystals?
 - □ yes, it can
- are boundary layer or free tropospheric aerosols more important?
 - free tropospheric
- is Hallett-Mossop ice multiplication important?
 - yes, but not to number concentration
- is heterogeneous ice nucleation important?
 - yes, at high concentrations

Future Work

- improve aerosol representation
 - WB-57 CN size distributions
 - Citation ice nuclei
 - chemical composition
- compare predicted updraft core particle size distributions with observations
- compare predicted cloud base and cloud top properties with observations
- estimate measurement uncertainty
- estimate model uncertainty
 - □ time
 - space
 - □ bin resolution